

---

## MATH 353: Engineering Mathematics III – Section 012

Spring 2013 (F.–J. Sayas)

Lab # 7

April 5

---

Open Matlab and *move to the Desktop or to a folder where you can find your work* at the end of the session. Type these lines

```
>> diary myworkApril5
>> format long
>> format compact
```

Download the function `heun.m` and the script `scriptApril5.m` from my website.

1. The function `heun.m` corresponds to the implementation of Heun's method (the explicit trapezoidal method) to numerically solve

$$y' = f(t, y) \quad a \leq t \leq b, \quad y(a) = y_a.$$

The script `scriptApril7.m` shows you one example of how this works. We are trying to solve the differential equation

$$(t^2 + 1)y' + 3ty = 6t, \quad y(0) = 1.$$

First of all, we need to write it in explicit form

$$y' = \frac{6t - 3ty}{t^2 + 1}, \quad y(0) = 1.$$

The exact solution of this problem is

$$y(t) = 2 - \frac{1}{(t^2 + 1)^{3/2}}.$$

- Run the script and figure out what we did. What is the interval where we solved the equation? How many time steps? Some place in the script, we have computed

$$E_h = \max_{0 \leq j \leq n} |w_j - y(t_j)|.$$

What is this value?

- Run the code again with  $n = 20$  time steps in the same time interval.
  - Run the code again with  $n = 100$  time steps in the interval  $[0, 10]$ .
2. Let us now compute errors for increasing values of  $n$ . Run the same example in the interval  $[0, 10]$ , with  $n = 10, 20, 40, 80, 160, 320$ , compute the errors, make a loglog plot of the errors and compare them with a loglog plot of  $(h, h^2)$ . To help you get organized...
    - Create a list `listn=[10 20 ...]`
    - Compute the vector with all values of  $h$  (you'll need it for the plots)

- Run `heun` when `n` takes values in the list `listn`
  - Compute the error and accumulate it on a vector of errors.
  - Do the loglog plots.
3. As we saw in class, there's an easy modification of this code that gives you the code for Euler's method. In this case the error is  $E_h = \mathcal{O}(h)$ , as opposed to  $E_h = \mathcal{O}(h^2)$  in Heun's method.
- Create the function `euler.m` following the same model as `heun.m` (copy-paste at will).
  - Repeat the experiment of Exercise 2 and show that you have order one and not two.
4. We are now going to experiment with a more complicated equation. For this one, we do not know the solution:

$$y' = y \cos t, \quad 0 \leq t \leq 8\pi \quad y(0) = 1.$$

Solve this for  $n = 25, 50, 100, 200, 400$  using Heun's method. At the point of getting the result I'm asking you to do the following:

- Compute the solution for  $n = 25$  and plot it. Pause (the command `pause` will wait for you to click on enter) and hold on.
  - Compute the solution for  $n = 50$  and plot it.
  - Go on until you have all experiments on the same graph.
5. Repeat Exercise 4 with Euler's method.